



Inbreeding in three-spined sticklebacks (*Gasterosteus aculeatus* L.): effects on testis and sperm traits

MARION MEHLIS^{1*}, JOACHIM G. FROMMEN^{1,2}, ANNA K. RAHN¹ and THEO C. M. BAKKER¹

¹*Institute for Evolutionary Biology and Ecology, University of Bonn, An der Immenburg 1, D-53121 Bonn, Germany*

²*Department of Behavioural Ecology, Institute of Ecology and Evolution, University of Bern, Wohlenstrasse 50a, CH-3032 Hinterkappelen, Switzerland*

Received 19 April 2012; revised 15 May 2012; accepted for publication 15 May 2012

Mating between relatives often results in inbreeding depression, and is assumed to have a strong effect on fitness traits such as fertility and gonad/gamete quality. However, data concerning this topic are contradictory and particularly scarce in fishes. Three-spined sticklebacks (*Gasterosteus aculeatus* L.) show inbreeding depression in fertilization and hatching success, survival rates, body symmetry and behavioural traits. To date, any knowledge of the impact of inbreeding on males' gonads and gametes is lacking in this species. In the present study, testis and sperm traits were quantified in outbred and inbred males. Overall, these traits were not generally impaired by inbreeding, and this result was not changed by a second/third generation of brother–sister matings. However, testes brightness, a potential measure of oxidative stress, was negatively correlated with sperm number. Additionally, inbred males with higher body condition had significantly brighter testes, whereas their sperm number was significantly negatively correlated with sperm quality (as estimated by head volume). Such a trade-off did not appear in outbred males. The comparatively small impact of inbreeding on testis and sperm traits might be explained by the low number of inbred individuals that reached the reproductive phase. © 2012 The Linnean Society of London, *Biological Journal of the Linnean Society*, 2012, 107, 510–520.

ADDITIONAL KEYWORDS: fish – heterozygosity – reproduction – sexual selection – sperm competition – testicular melanization.

INTRODUCTION

Mating with close relatives leads to an increased level of homozygosity (Charlesworth & Charlesworth, 1987), resulting in a loss of fitness in many species (Crnokrak & Roff, 1999; Armbruster & Reed, 2005). This inbreeding depression might influence many different traits. For example, inbred individuals face a reduced resistance to parasites (Coltman *et al.*, 1999) and are more vulnerable to environmental changes (Keller *et al.*, 1994). Furthermore, inbreeding may lead to a higher degree of body deformations, resulting in a higher mortality rate (Waldman & McKinnon, 1993). However, sometimes mating with close relatives has

no detrimental effects (Keane, Creel & Waser, 1996; Sheridan & Pomiankowski, 1997; Peer & Taborsky, 2005; Thünken *et al.*, 2007) or even increases an individual's fitness, as an individual that mates with a relative will enable that relative to spread genes identical by descent (Kokko & Ots, 2006).

Traits closely related to fitness (such as fertility or survival) are known to be more strongly influenced by inbreeding depression than traits that are less closely related to fitness (Falconer, 1989; DeRose & Roff, 1999). This difference might be explained by larger directional dominance in fitness-related traits (DeRose & Roff, 1999) or by the fact that, on average, fitness-related traits are influenced by more loci (Houle, 1992; Keller & Waller, 2002). Thus, fertility as well as gonadal and gamete traits are expected to suffer strong consequences of inbreeding. However, studies investigating such a connection have yielded

*Corresponding author.

E-mail: mmehlis@evolution.uni-bonn.de

contradictory results. For example, a study on Shetland ponies (*Equus ferus caballus* L.) found decreasing fertility with increasing homozygosity (van Eldik *et al.*, 2006), whereas a study on Austrian Noriker draught horses did not find such a negative effect (Aurich, Achmann & Aurich, 2003). Margulis & Walsh (2002) demonstrated that testes mass of Alabama beach mice *Peromyscus polionotus* (Wagner) decreased when inbreeding coefficients increased. Similarly, in wild rabbits *Oryctolagus cuniculus* (L.), the production of abnormal sperm was negatively correlated with heterozygosity (Gage *et al.*, 2006). Furthermore, in some species, the negative impact of inbreeding on fertility is limited to only one sex, whereas the other sex remained unaffected (Brekke *et al.*, 2010; Rioux-Paquette, Festa-Bianchet & Coltman, 2011). For example, in the butterfly *Bicyclus anynana* (Butler), approximately one-half of the male offspring of brother–sister matings was infertile, whereas female fertility was unaffected by inbreeding (Saccheri *et al.*, 2005). Also, in a small Swiss human population, inbred women had fewer children, whereas the reproductive success of men was not affected (Postma, Martini & Martini, 2010).

Fluctuating asymmetry is known to be a reliable indicator of developmental and environmental stress (Leary & Allendorf, 1989; Sopinka *et al.*, 2012). Also, genetic stress as a result of inbreeding could lead to fluctuating asymmetry. For example, the degree of fluctuating asymmetry was positively correlated with the inbreeding coefficient and negatively correlated with the proportion of normal sperm in the Cuvier's gazelle *Gazella cuvieri* (Ogilby) (Roldan *et al.*, 1998). A comparative study in three more species of ungulates showed that fluctuating asymmetry is related to individuals semen quality (Gomendio, Cassinello & Roldan, 2000). However, fluctuating testis asymmetries have seldom been investigated. Furthermore, the impact of inbreeding on males' gonadal and sperm traits has been less well explored, especially in fish (but see Zajitschek & Brooks, 2010). Thus, the present study aimed to increase the knowledge of the effects of inbreeding on males' reproductive traits.

The three-spined stickleback (*Gasterosteus aculeatus* L.) is an ideal model species for studying the influence of inbreeding on testis and sperm traits for several reasons. First, anadromous sticklebacks frequently colonize new fresh water environments and establish new populations consisting of only a few founder individuals, hence representing a rather small gene pool (Raeymaekers *et al.*, 2005; Ólafsdóttir, Snorrason & Ritchie, 2007; Wund *et al.*, 2008). The risk of inbreeding and associated inbreeding depression in such newly-established populations should be high (Aeschlimann *et al.*, 2003). Indeed, the

three-spined stickleback show inbreeding depression in life history traits such as fertilization success, hatching rates, and mortality of both juveniles and adults (Frommen *et al.*, 2008), as well as in behavioural traits (Frommen *et al.*, 2007) and body symmetry (Mazzi *et al.*, 2004). Inbreeding depression in this species can be caused by just one generation of brother–sister mating, and is further intensified by a second generation of incestuous mating (Frommen *et al.*, 2008). However, several traits under sexual selection such as mating decisions (Frommen & Bakker, 2006; Mehliis, Bakker & Frommen, 2008; but see also Mazzi *et al.*, 2004), aggressive interactions (Mehliis *et al.*, 2009) or the expression of male breeding coloration (Frommen *et al.*, 2008), appeared not to be affected by inbreeding.

Second, the stealing of fertilizations, so-called 'sneaking', is a common behavioural tactic in this species (Goldschmidt, Foster & Sevenster, 1992; Jamieson & Colgan, 1992; Rico, Kuhnlein & Fitzgerald, 1992; Largiadèr, Fries & Bakker, 2001; Vliieger & Candolin, 2009), which consequently results in sperm competition (Parker, 1990). Spermatogenesis in sticklebacks takes place in the months in which photoperiod is short (Borg, 1982). Thus, sperm are not produced during the breeding season, which takes place between April and August, resulting in a larger number of sperm stored in testes of virgin males (e.g. early spring) compared to those of multiply-mated males (e.g. late summer) (Zbinden, Largiadèr & Bakker, 2001). As the sperm quantity decreases during the course of the breeding season, sperm are allocated carefully (Zbinden *et al.*, 2003; Zbinden, Largiadèr & Bakker, 2004). Furthermore, gonadal mass is related to reproductive success (Cubillos & Guderley, 2000), and the number of stored sperm is significantly positively correlated with the gonadosomatic index (Zbinden *et al.*, 2001). Thus, a stickleback male that is unable to produce sufficient amounts of fertile sperm because of inbreeding depression will most probably suffer a reduced fertilization success.

Using the same breeding line as that described by Frommen *et al.* (2008), the present study investigated how inbreeding affects testis and sperm traits of the three-spined stickleback. We measured the potential impact of inbreeding on the gonadosomatic index, sperm number, and sperm morphology. Furthermore, we analyzed melanophore pigmentation of the testes, which might play a role in protection against ultraviolet (UV)-light and/or oxidative stress (Kaidbey *et al.*, 1979; Plonka *et al.*, 2009; Galván, Møller & Erritzøe, 2011). Finally, asymmetries of testis traits were considered as a measure of developmental and genetic stress (Leary & Allendorf, 1989; Mazzi, Largiadèr & Bakker, 2002). All measures were

compared between males that were outbred or inbred for one or several generations.

MATERIAL AND METHODS

EXPERIMENTAL SUBJECTS

Three-spined sticklebacks from an anadromous population were caught during their spring migration in April 2002, 2003, and 2004 on the island of Texel, the Netherlands. During this time, large numbers of sticklebacks migrate from the ocean into freshwater habitats (Kemper, 1995). Furthermore, individuals captured on Texel are representative of the whole population, and the gene pool is highly heterogeneous (Heckel *et al.*, 2002).

Fish used in the present study were outbred and first-, second- or third-generation inbred descendants of these wild-caught parents. Outbred fish were the F_1 progeny of randomly crossed wild-caught fish. To achieve different outbred families (full-sib groups), males were allowed to spawn with one single randomly chosen female. Inbred fish were then produced by three subsequent generations of brother–sister matings. Thus, first-generation inbred fish were the F_2 progeny, second-generation inbred fish were the F_3 progeny, and third-generation inbred fish were the F_4 progeny of the wild-caught fish. To avoid pseudoreplication, inbred clutches were obtained by using only one parental pair for each family. Outbred clutches were produced in April and May 2004, whereas inbred clutches were produced in December 2004 and January 2005. To exclude paternal and tank effects, all clutches were removed from the males' tank and split into two groups 1 h after fertilization (Frommen *et al.*, 2008). Fish were kept in full-sib groups of 10–20 individuals. The holding tanks measured 50 cm × 30 cm × 30 cm (length × width × height) and were placed in an air-conditioned room under standardized winter light-regime (8 : 16 h light/dark cycle at 17 ± 1 °C). To simulate the start of the breeding season, the light regime was changed to summer conditions in August 2006 (16 : 8 h light/dark cycle at 17 ± 1 °C). Males that showed initial signs of breeding coloration in the holding tanks were isolated in separate tanks (40.5 × 20.5 × 25 cm) for nest building (Mehlis *et al.*, 2008). All males used were virgin because they had no possibility to build a nest in their holding tank and were not paired with any female in their nesting tank. In addition, only males that had built a nest were used, which ensured that they were reproductively active. On average, males needed one day (median and quartiles: 1, 1, and 2) to build their nest. They were allowed to care for the nest for up to 18 days (median and quartiles: 10, 8, and 18). Before they were quickly sacrificed, males' standard length

(SL) and body mass (M) were measured, to calculate their body condition (BC) *sensu* Bolger & Connolly (1989):

$$BC = 100 \times M/SL^3 \quad (1)$$

In total, 125 males were used originating from 31 outbred families (a total of 85 males) and 15 inbred families (a total of 40 males). Of the inbred families seven families (18 males) were inbred for one generation, six families (18 males) were inbred for two generations and two families (four males) were inbred for three generations. Males from outbred and inbred families did not differ significantly with respect to the duration of nest building, in the time they cared for a nest, in SL, in M or in BC ($N_{\text{outbred}} = 31$, $N_{\text{inbred}} = 15$; unpaired Wilcoxon-test, all $W < 291$, all $P > 0.173$; see Supporting information, Table S1).

MEASUREMENT OF TESTIS TRAITS

Before testes dissection (April 2008 and June 2009), all males had been stored in 70 % ethanol for at least 2 years, which might have led to dehydration and hence organ deformation. Thus, new body measurements of males were taken using the same protocol as before and the new measurements were used for all further analyses. Afterwards, both testes were removed and placed separately on a tissue to remove adhered liquids. Then they were weighed to the nearest milligram (OHAUS, Explorer, E11140) and stored separately in Eppendorf tubes containing 500 µl of 70 % ethanol. The gonadosomatic index (GSI) was calculated *sensu* de Vlaming, Grossman & Chapman (1982):

$$GSI = (TM M^{-1}) \times 100 \quad (2)$$

where TM is the sum of the mass of both testes ($TM_{\text{left}} + TM_{\text{right}}$).

To determine the intensity of melanophore pigmentation, a standardized image of each testis was taken with a digital camera (HV-C20AMP; Hitachi Denshi Ltd) mounted on a binocular (Leica S8AP0) using Diskus 4.6 (C.H. Hilgers Technical office). To ensure comparable light regimes, all specimens were illuminated using a cold light source (KL 1500 LCD Leica), which was always placed in the same position. Furthermore, the orientation of the testis was always the same with the outside upturned. (Fig. 1A, B, C). Pictures were analyzed in ADOBE PHOTOSHOP CS4 (Adobe Systems Inc.) using the CIE $L^*a^*b^*$ colour space, which is a method that is frequently used in fish colour evaluation (Adachi *et al.*, 2005; Svensson *et al.*, 2005). A white Munsell card (N10) visible on each picture served as white balance. To obtain the




Figure 1. Testes and sperm of three-spined stickleback. A, B, C, variation in testis melanophore pigmentation and the corresponding mean brightness (L^* value; A1–C1). Munsell chips are visible on the left side and a millimetre scale is provided at the bottom of each image. D, sperm morphology traits, the dotted line indicates flagellum length and the solid line indicates head length.

mean L^* value [brightness; ranging from black (0) to white (100)] of each testis, testis shape was marked using the magnetic-lasso-tool. The testes of three-spined sticklebacks show a high variability in melanophore pigmentation (Fig. 1A, B, C) and the L^* value (brightness) is a good indicator of melanophore density (Fig. 1A1, B1, C1). Brightness was determined using the entire area inside of the magnetic-

lasso-tool for each male's testis (L^*_{left} and L^*_{right}) and averaged (L^*_{total}). Finally, asymmetries in testis mass (Eqn 3) and testis brightness (Eqn 4) were calculated:

$$\text{Testis mass asymmetry} = (\text{TM}_{\text{heavy}} - \text{TM}_{\text{light}}) / \text{TM}_{\text{heavy}} \quad (3)$$

$$\text{Brightness asymmetry} = (L^*_{\text{bright}} - L^*_{\text{dark}}) / L^*_{\text{bright}} \quad (4)$$

MEASUREMENTS OF SPERM TRAITS

The number of sperm stored in left and right testis are highly correlated in three-spined sticklebacks of the study population (Bakker *et al.*, 2006). Therefore, the measurement of sperm number was solely conducted on the left testis. To obtain sperm, each male's left testis was pestled in 200 μl of phosphate-buffered saline (Dulbecco & Vogt, 1954). Ten microlitres of this solution were diluted with 190 μl of tap water to reduce sperm density. Twelve microlitres of this dilution were filled into a Neubauer improved counting chamber (Labor Optik, 0.0025 mm^2 , depth 0.1 mm). Sperm were counted in 64 cells, which were equally distributed over the chamber. Thus, the total number of sperm (S_T) was calculated using the equation:

$$S_T = S_A \times 4000 \times 200 \times 20 \quad (5)$$

where S_A is the mean sperm number of the 64 counted cell chambers.

Sperm morphology was determined by scanning electron microscopy (SEM; Fig. 1D) with a subsample of 23 males from different families [twelve outbred and eleven inbred males (four males first-, six males second- and one male third-generation inbred, which were pooled for analyses)]. The preparation of sperm for SEM was carried out *sensu* Mortimer (1994) and was adapted for stickleback sperm (see Supporting information, Appendix S1). Sperm variables were measured using 2500–4500 magnified SEM-images (digitized Leitz AMR 1000). They were digitized using DISS (Digital Image Scanning System, Point Electronic GmbH) and DIPS (Digital Image Processing System, version 2.5.2.1, Point Electronic GmbH). Although ethanol has been used as a preservative medium in a number of comparative studies concerning testes and sperm variables (Byrne, Roberts & Simmons, 2002; Kvarnemo & Simmons, 2004), sperm midpiece and head could not visually be distinguished as a result of the dehydrating effects of the alcohol and were thus measured together. However, because both inbred and outbred males had been stored in 70 % ethanol for the same time, data were comparable with each other. Head length (hl) and flagellum length (fl) were measured only on images that showed intact sperm with a well visible head and a complete tail (Fig. 1D). In total, 187 sperm were measured

(median and quartiles: 6, 4, 12.5 per male) using IMAGEJ (NIH). The measures used are known to be good proxies of reproductive success in several animal species (Murray, Kozłowska & Cutter, 2011), including fish (Boschetto, Gasparini & Pilastro, 2011). In addition, several studies revealed that sperm morphology is linked to sperm velocity (Fitzpatrick, Garcia-Gonzalez & Evans, 2010). Thus, as a proxy of sperm motility, the head length to flagellum length ratio (Eqn 6) (Humphries, Evans & Simmons, 2008) and the complete head volume (including the mid-piece volume; Anderson & Dixson, 2002) (Eqn 7) assuming the form of a sphere (Fig. 1D) were calculated:

$$\text{Head to flagellum ratio} = \text{hl/fl} \quad (6)$$

$$\text{Head volume} = \frac{1}{6} \times \pi \times (\text{hl})^3 \quad (7)$$

The head to flagellum ratio is a good measure of sperm motility in three-spined sticklebacks, and flagellum length, mid-piece volume and head volume were shown to predict fertilization success in this species (Marion Mehlis, Lukas K. Hilke, Theo C. M. Bakker unpubl. data and Theo C. M. Bakker, Michael Hollmann, Marion Mehlis, Marc Zbinden, unpubl. data).

REPEATABILITY OF TESTIS AND SPERM TRAITS

To calculate the repeatability (R) and its standard error (SE; after Becker, 1992) of the measurements of testis traits, one male was randomly chosen from each family (approximately 37 % of total sample size). From these males, TM_{right} and $\text{L}^*_{\text{right}}$ were measured three times. Furthermore, from the subsample of 23 males, one randomly chosen sperm was measured twice from each male (approximately 12 % of total sample size). The formula used to calculate repeatability was:

$$R = \left(\frac{[(MS_a - MS_w)/N_0]}{[(MS_a - MS_w)/N_0] + MS_w} \right) \times 100 \quad (8)$$

where MS_a is the between sperm mean square, MS_w is the within sperm mean square, and N_0 equals three (testis traits) and two (sperm traits) respectively. Repeatability was high for the measurement of testis mass (89.71 ± 0.15 %), as well as for testis brightness (98.91 ± 0.02 %). The repeatability values were also high for the measurement of sperm traits. This was true for head length (94.24 ± 0.35 %), as well as for flagellum length (99.92 ± 0.01 %).

STATISTICAL ANALYSIS

The R 2.9.1 statistical package was used for analyses. The mean values of the different families were used to avoid pseudoreplication. Thus, the sample size for analyses of testis traits was $N_{\text{outbred}} = 31$ and $N_{\text{inbred}} =$

15 ($N_{\text{1st inbred}} = 7$, $N_{\text{2nd + 3rd inbred}} = 8$) and, for analyses of sperm traits, $N_{\text{outbred}} = 12$ and $N_{\text{inbred}} = 11$. Second- and third-generation inbred families were pooled because of the low sample size ($N_{\text{3rd inbred}} = 2$). For statistical analyses, linear models ('lm') were conducted using the respective testis or sperm traits as dependent variables and the breeding regime (outbred versus inbred and outbred versus various levels of inbreeding (first- and second + third-generation inbred) as an explanatory variable (for a detailed list of relevant models, see the Supporting information, Table S2). Inbred males were approximately 8 months younger than outbred ones. Fish grow throughout their life and, in most cases, reproductive success is positively related with size (Reznick, Ghalambor & Nunney, 2002), which could also be shown for three-spined sticklebacks (Allen & Wootton, 1982; Kraak, Bakker & Mundwiler, 1999). Similarly, in the present study, age was significantly correlated with male SL (Spearman correlation: $N = 125$; $r_s = 0.206$, $P = 0.021$). Thus, as a surrogate for age, SL was included as an explanatory variable in all models and never removed to control for possible age influences. To check for differences in the relationship among testis and sperm investment between outbred and inbred males, additional linear models ('lm') were conducted, in which interactions between the breeding regime and different testis respectively sperm traits were included (for a detailed list of relevant models, see Supporting information, Table S2). For all models, tests of significance were based on likelihood-ratio tests (LRT). Because normally distributed dependent variables are a prerequisite for running linear models, sperm number was square-root transformed, GSI was inverse transformed, and head volume and asymmetries in testis mass and testis brightness were logarithmically transformed, to achieve normal distributions of the residuals of the best explaining model, according to Kolmogorov–Smirnov tests. All P -values are based on two-tailed tests.


RESULTS

TESTIS TRAITS

Males showed considerable variation in GSI and testis brightness ($\text{L}^*_{\text{total}}$) (Fig. 1, Table 1), which did not differ significantly between outbred and inbred males ($N_{\text{outbred}} = 31$, $N_{\text{inbred}} = 15$; 'lm', both $\chi^2 < 0.082$, both $P > 0.776$; Table 1). Differences in GSI and testis brightness did not differ significantly between different levels of inbreeding ($N_{\text{outbred}} = 31$, $N_{\text{1st inbred}} = 7$, $N_{\text{2nd + 3rd inbred}} = 8$; 'lm', both $\chi^2 < 0.913$, both $P > 0.409$; Table 1). Asymmetries of testis mass or brightness were not significantly different between outbred and inbred males ($N_{\text{outbred}} = 31$, $N_{\text{inbred}} = 15$; 'lm', both $\chi^2 < 1.215$, both $P > 0.277$; Table 1). The level

Table 1. Descriptive statistics (median, first and third quartile) of testis and sperm traits used in the analyses

	Outbred			Inbred			First-generation inbred			Second + third-generation inbred		
	Median	First quartile	Third quartile	Median	First quartile	Third quartile	Median	First quartile	Third quartile	Median	First quartile	Third quartile
Gonadosomatic index	0.52	0.44	0.71	0.48	0.43	0.60	0.52	0.41	0.75	0.46	0.43	0.55
Testis brightness (L^*_{total})	49.50	41.44	51.70	50.63	43.10	52.91	42.00	41.29	48.11	52.38	49.34	54.98
Testis mass asymmetry	0.210	0.167	0.337	0.239	0.139	0.265	0.227	0.138	0.253	0.255	0.195	0.282
Testis brightness asymmetry	0.092	0.055	0.124	0.098	0.074	0.156	0.118	0.075	0.156	0.096	0.074	0.157
Sperm number	12.6×10^6	8.9×10^6	14.8×10^6	11.9×10^6	8.9×10^6	16.5×10^6	16.3×10^6	12.5×10^6	17.0×10^6	9.9×10^6	7.8×10^6	12.1×10^6
Flagellum length (μm)	18.19	16.37	19.11	16.84	15.84	21.21	-	-	-	-	-	-
Head length (μm)	0.72	0.65	0.79	0.69	0.67	0.77	-	-	-	-	-	-
Head to flagellum ratio	0.042	0.038	0.045	0.039	0.035	0.044	-	-	-	-	-	-
Head volume (μm^3)	0.217	0.155	0.280	0.197	0.165	0.272	-	-	-	-	-	-

**Figure 2.** Significant interactions between outbred (circles; solid lines) and inbred (crosses; dotted lines) males. A, relationship between male body condition and testis brightness (L^*_{total}). B, relationship between sperm number and head volume (μm^3).

of inbreeding was not significantly related to asymmetries of testis mass or brightness ($N_{\text{outbred}} = 31$, $N_{\text{1st inbred}} = 7$, $N_{\text{2nd + 3rd inbred}} = 8$; 'lm', both $\chi^2 < 0.848$, both $P > 0.436$; Table 1).

SPERM TRAITS

Sperm number did not differ significantly between outbred and inbred males ($N_{\text{outbred}} = 31$, $N_{\text{inbred}} = 15$; 'lm', $\chi^2 = 0.417$, $P = 0.522$; Fig. 2, Table 1). Analysing the different generations of inbreeding separately revealed no significant differences in number of sperm with increasing levels of inbreeding ($N_{\text{outbred}} = 31$, $N_{\text{1st inbred}} = 7$, $N_{\text{2nd + 3rd inbred}} = 8$; 'lm', $\chi^2 = 2.021$, $P = 0.145$; Table 1). Furthermore, outbred and inbred males did not differ significantly in sperm morphology traits, neither in head length, flagellum length, head to flagellum ratio, nor in head volume ($N_{\text{outbred}} = 12$, $N_{\text{inbred}} = 11$; 'lm', all $\chi^2 < 0.718$, all $P > 0.407$; Table 1).

INTERACTIONS BETWEEN TESTIS AND SPERM TRAITS

The number of sperm was negatively correlated with testis brightness (L^*_{left}), which was true for

inbred (Spearman correlation: $N_{\text{inbred}} = 15$; $r_s = -0.654$, $P = 0.010$), as well as for outbred males (Spearman correlation: $N_{\text{outbred}} = 31$; $r_s = -0.371$, $P = 0.040$). This means that, in all males, brighter testes contained less sperm. Furthermore, inbred males in better BC had significantly brighter testes (L^*_{total}) (Pearson correlation: $N_{\text{inbred}} = 15$; $r_p = 0.640$, $P = 0.010$; Fig. 2A), whereas this relationship was not found in outbred males (Pearson correlation: $N_{\text{outbred}} = 31$; $r_p = -0.191$, $P = 0.304$; Fig. 2A), leading to a significant interaction between outbred and inbred males with respect to the relationship between testis brightness and BC ($N_{\text{outbred}} = 31$, $N_{\text{inbred}} = 15$; 'lm', $\chi^2 = 4.676$, $P = 0.037$). Furthermore, the relationship between sperm number and head volume differed significantly between outbred and inbred males ($N_{\text{outbred}} = 12$, $N_{\text{inbred}} = 11$; 'lm', $\chi^2 = 6.054$, $P = 0.024$). Inbred males with higher numbers of sperm had significantly smaller head volumes (Spearman correlation: $N_{\text{inbred}} = 11$; $r_s = -0.673$, $P = 0.028$; Fig. 2B), whereas the opposite was true for outbred males (Pearson correlation: $N_{\text{outbred}} = 12$; $r_p = 0.596$, $P = 0.041$; Fig. 2B).

DISCUSSION

Outbred and inbred males did not differ in GSI, testis coloration, degree of gonadal asymmetries, sperm number, and sperm morphology. As discussed by Keller & Waller (2002), it might be difficult to detect the effects of inbreeding when inbreeding depression is high. It is argued that, in such a case, homozygous individuals would suffer from lethal mutations early in development and might not reach the adult stage. This could lead to low estimates of the effects of inbreeding, as shown for example in the Scots pine (*Pinus sylvestris* L.) (Savolainen & Hedrick, 1996). The present study used the same breeding line as that employed in Frommen *et al.* (2008). Frommen *et al.* (2008) showed that sticklebacks are adversely affected by inbreeding depression at an early life stage because fertilization and hatching rates in inbred offspring were low and the mortality of juveniles was high. It is therefore possible that inbred fish, which had suffered strongly from inbreeding depression, died before they reached the reproductive phase and were not measured in the present study. Thus, inbred fish that do reach adulthood may not suffer as much from inbreeding depression and their sperm and gonadal traits may be of similar quality to those of outbred conspecifics. Furthermore, in the present study, both outbred and inbred males were laboratory-bred. Standardized near-optimal laboratory rearing conditions could have overruled the possible negative effects of inbreeding. In the wild, diseases and restricted food availability may have detrimental effects on males, so that they may not be

able to show comparable fertility and hence fitness with respect to outbred males. For example, in male house mice (*Mus musculus* L.), reproductive success was more strongly influenced by inbreeding when mice were reared in large semi-natural enclosures (Meagher, Penn & Potts, 2000), implying that competition in the wild might increase the harmful effects of inbreeding.

A further reason for a lack of inbreeding depression in testis and sperm traits could be that the applied levels of inbreeding did not reduce heterozygosity sufficiently strongly to unmask deleterious recessive alleles. Indeed, several studies found that inbreeding depression was intensified by on-going incestuous matings (Richards, 2000; Crnokrak & Barrett, 2002; Dolgin *et al.*, 2007). The results of Frommen *et al.* (2008) support this idea as well. One generation of inbreeding by brother–sister matings significantly lowered the fertilization success and hatching rate of eggs. This effect was intensified by a second generation of inbreeding (Frommen *et al.*, 2008). In the present study, the increase of inbreeding depression by a second generation of incestuous mating was marginal if anything. GSI, testis brightness, and asymmetries, as well as sperm number and morphology, did not differ significantly between outbred, one generation inbred and second + third-generation inbred males.

In several vertebrates, testes are covered by melanophores that contain melanin, such as in fish (Louiz, Ben-Attia & Ben-Hassine, 2009), amphibians (Zieri, Taboga & De Oliveira, 2007), reptiles (Guillette, Weigel & Flater, 1983), birds (Galván *et al.*, 2011), and mammals (Poole & Lawton, 2009). Melanophores contain dark pigments, which are assumed to protect the testes from deleterious UV light as a result of their strong light absorbance (Kaidbey *et al.*, 1979; Plonka *et al.*, 2009) or to protect male germ cells from oxidative stress (Galván *et al.*, 2011). In birds, for example, testis melanization has evolved in species with high rates of accumulated mitochondrial mutations and is assumed to be an adaptive response related to the protective capacity of melanin against oxidative stress (Galván *et al.*, 2011). In the present study, a higher number of melanophores was related to a higher number of sperm, both in outbred and inbred males. Thus, males that were of generally higher quality in terms of sperm production were also of higher quality in terms of sperm protection. However, testes melanization in inbred males was negatively correlated with BC, which means that males with higher BC had testes with fewer melanophores. Thus, inbred males might face a trade-off between investment in BC and testis melanization: the higher their investment in BC the smaller was the investment into melanophores. Because sperm

are known to be highly prone to suffer from oxidative stress (Aitken & Baker, 2006), inbred sticklebacks may not be able to keep a high BC and protect their sperm from oxidative stress at the same time. By contrast, such a trade-off was not found in outbred fish, and might thus be interpreted as a cryptic case of inbreeding depression. Additionally, inbred males (in contrast to outbred males) faced a trade-off between sperm quantity and sperm quality (as estimated by head volume). Hence, inbred males might either be able to invest into producing larger ejaculates, or in smaller ejaculates containing better sperm. Taken together, these results show that, for inbred males, it is physiologically taxing to produce large numbers of sperm of high quality, and protect them from potential damage caused by UV light and oxidative stress, at the same time as maintaining a high BC. In nature, sticklebacks face a high risk of sperm competition (Goldschmidt *et al.*, 1992; Jamieson & Colgan, 1992; Largiadèr *et al.*, 2001). Under these conditions, it is advantageous for an individual to possess both a large amount of sperm and high-quality sperm. Furthermore, stickleback males in good BC are more successful in attracting mates and withstanding competitors (Bakker & Milinski, 1993). According to the present results, inbred males should face problems in both of these terms.

Outbred and inbred males were of different age, with the inbred fish being, on average, 8 months younger than the outbred fish. Hence, an influence of age on the experimental results cannot be completely ruled out. For example, other studies found a negative effect of age on testis and sperm traits even in three-spined sticklebacks (M. Mehlis & T. C. M. Bakker, unpubl. data; de Fraipont, FitzGerald & Guderley, 1993). However, these studies compared stickleback males that had experienced one breeding season with older males that had experienced two breeding seasons. In the present study, both groups were in their first reproductive season when measurements were taken, and were exposed to the same duration of winter conditions before the light-regime was switched to resemble summer again. Borg (1982) showed that, in three-spined sticklebacks, spermatogenesis was only active during short photoperiods. The time span for sperm production was the same for both outbred and inbred males, indicating that stored sperm were of similar age in both groups. In addition, despite the age difference between outbred and inbred males, they did not differ significantly in body measurements, supporting the comparability of the two groups. Moreover, we added SL as an additional variable to our models, adjusting for nonsignificant differences in body size between the groups.

In conclusion, the present study shows that sperm and gonadal traits are not generally impaired by

one or more generations of brother-sister matings. However, inbred males appeared to face rather cryptic inbreeding depression because they were unable to produce large numbers of sperm, adequately protected against potential damage caused by UV light or oxidative stress, and, at the same time, maintain a good BC. Finally, inbred fish were unable to both produce large amounts of sperm and maintain high-quality sperm. Such a trade-off did not manifest in outbred males, which might imply that inbred males were of lower quality and may have a fitness disadvantage, especially when confronted with sperm competition under natural conditions.

ACKNOWLEDGEMENTS

We are grateful to the 'Bakker' research-group, especially Ingolf P. Rick, for discussions. Meike Hiermes and Valentina Balzarini are acknowledged for correcting our English. We thank Corinna Luz for help in rearing sticklebacks and Jan Hottentot for catching sticklebacks. Stefanie Gierszewski, Anh-Phong Nguyen, Dagmar Wenzel and Erika Müller-Schulte are acknowledged for their help with measuring the SEM images. This research was funded by the Deutsche Forschungsgemeinschaft (DFG) (BA 2885/4-1).

REFERENCES

- Adachi K, Kato K, Wakamatsu K, Ito S, Ishimaru K, Hirata T, Murata O, Kumai H. 2005. The histological analysis, colorimetric evaluation, and chemical quantification of melanin content in 'suntanned' fish. *Pigment Cell Research* **18**: 465–468.
- Aeschlimann PB, Häberli MA, Reusch TBH, Boehm T, Milinski M. 2003. Female sticklebacks *Gasterosteus aculeatus* use self-reference to optimize MHC allele number during mate selection. *Behavioral Ecology and Sociobiology* **54**: 119–126.
- Aitken RJ, Baker M. 2006. Oxidative stress, sperm survival and fertility control. *Molecular and Cellular Endocrinology* **250**: 66–69.
- Allen JRM, Wootton RJ. 1982. Age, growth and rate of food consumption in an upland population of the three-spined stickleback, *Gasterosteus aculeatus* L. *Journal of Fish Biology* **21**: 95–105.
- Anderson MJ, Dixon AF. 2002. Sperm competition: motility and the midpiece in primates. *Nature* **416**: 496.
- Armbruster P, Reed DH. 2005. Inbreeding depression in benign and stressful environments. *Heredity* **95**: 235–242.
- Aurich C, Achmann R, Aurich J. 2003. Semen parameters and level of microsatellite heterozygosity in Noriker draught horse stallions. *Theriogenology* **60**: 371–378.
- Bakker TCM, Milinski M. 1993. The advantages of being red: sexual selection in the stickleback. *Marine Behaviour and Physiology* **23**: 287–300.

- Bakker TCM, Zbinden M, Frommen JG, Weiss A, Largiadèr CR. 2006.** Slow fertilization of stickleback eggs: the result of sexual conflict? *BMC Ecology* **6**: 7.
- Becker WA. 1992.** *Manual of quantitative genetics*. Washington, DC: Academic Enterprises.
- Bolger T, Connolly PL. 1989.** The selection of suitable indices for the measurement and analysis of fish condition. *Journal of Fish Biology* **34**: 171–182.
- Borg B. 1982.** Seasonal effects of photoperiod and temperature on spermatogenesis and male secondary sexual characters in the three-spined stickleback, *Gasterosteus aculeatus*. *Canadian Journal of Zoology* **60**: 3377–3386.
- Boschetto C, Gasparini C, Pilastro A. 2011.** Sperm number and velocity affect sperm competition success in the guppy (*Poecilia reticulata*). *Behavioral Ecology and Sociobiology* **65**: 813–821.
- Brekke P, Bennett P, Wang J, Pettorelli N, Ewen J. 2010.** Sensitive males: inbreeding depression in an endangered bird. *Proceedings of the Royal Society of London Series B, Biological Sciences* **277**: 3677–3684.
- Byrne PG, Roberts JD, Simmons LW. 2002.** Sperm competition selects for increased testes mass in Australian frogs. *Journal of Evolutionary Biology* **15**: 347–355.
- Charlesworth D, Charlesworth B. 1987.** Inbreeding depression and its evolutionary consequences. *Annual Review of Ecology and Systematics* **18**: 237–268.
- Coltman DW, Pilkington JG, Smith JA, Pemberton JM. 1999.** Parasite-mediated selection against inbred Soay sheep in a free-living, island population. *Evolution* **53**: 1259–1267.
- Crnokrak P, Barrett SCH. 2002.** Perspective: purging the genetic load: a review of the experimental evidence. *Evolution* **56**: 2347–2358.
- Crnokrak P, Roff DA. 1999.** Inbreeding depression in the wild. *Heredity* **83**: 260–270.
- Cubillos ER, Guderley HE. 2000.** Analysis of the factors related with mate choice and reproductive success in male three-spined sticklebacks. *Journal of Fish Biology* **56**: 1201–1216.
- DeRose MA, Roff DA. 1999.** A comparison of inbreeding depression in life-history and morphological traits in animals. *Evolution* **53**: 1288–1292.
- Dolgin E, Charlesworth B, Baird S, Cutter A. 2007.** Inbreeding and outbreeding depression in *Caenorhabditis* nematodes. *Evolution* **61**: 1339–1352.
- Dulbecco R, Vogt M. 1954.** Plaque formation and isolation of pure lines with poliomyelitis viruses. *Journal of Experimental Medicine* **99**: 167–182.
- van Eldik P, van der Waaij E, Ducro B, Kooper A, Stout T, Colenbrander B. 2006.** Possible negative effects of inbreeding on semen quality in Shetland pony stallions. *Theriogenology* **65**: 1159–1170.
- Falconer DS. 1989.** *An introduction to quantitative genetic*. London: Wiley.
- Fitzpatrick JL, Garcia-Gonzalez F, Evans JP. 2010.** Linking sperm length and velocity: the importance of intramale variation. *Biology Letters* **6**: 797–799.
- de Fraipont M, FitzGerald GJ, Guderley H. 1993.** Age-related differences in reproductive tactics in the three-spined stickleback, *Gasterosteus aculeatus*. *Animal Behaviour* **46**: 961–968.
- Frommen JG, Bakker TCM. 2006.** Inbreeding avoidance through non-random mating in sticklebacks. *Biology Letters* **2**: 232–235.
- Frommen JG, Luz C, Mazzi D, Bakker TCM. 2008.** Inbreeding depression affects fertilization success and survival but not breeding coloration in threespine sticklebacks. *Behaviour* **145**: 425–441.
- Frommen JG, Mehlis M, Brendler C, Bakker TCM. 2007.** Shoaling decisions in three-spined sticklebacks (*Gasterosteus aculeatus*) – familiarity, kinship and inbreeding. *Behavioral Ecology and Sociobiology* **61**: 533–539.
- Gage M, Surridge A, Tomkins J, Green E, Wiskin L, Bell D. 2006.** Reduced heterozygosity depresses sperm quality in wild rabbits, *Oryctolagus cuniculus*. *Current Biology* **16**: 612–617.
- Galván I, Møller AP, Erritzøe J. 2011.** Testicular melanization has evolved in birds with high mtDNA mutation rates. *Journal of Evolutionary Biology* **24**: 988–998.
- Goldschmidt T, Foster SA, Sevenster P. 1992.** Inter-nest distance and sneaking in the three-spined stickleback. *Animal Behaviour* **44**: 793–795.
- Gomendio M, Cassinello J, Roldan ERS. 2000.** A comparative study of ejaculate traits in three endangered ungulates with different levels of inbreeding: fluctuating asymmetry as an indicator of reproductive and genetic stress. *Proceedings of the Royal Society of London Series B, Biological Sciences* **267**: 875–882.
- Guillette L, Weigel J, Flater G. 1983.** Unilateral testicular pigmentation in the Mexican lizard *Sceloporus variabilis*. *Copeia* **1983**: 155–161.
- Heckel G, Zbinden M, Mazzi D, Kohler A, Reckeweg G, Bakker TCM, Largiadèr CR. 2002.** Microsatellite markers for the three-spined stickleback (*Gasterosteus aculeatus* L.) and their applicability in a freshwater and an anadromous population. *Conservation Genetics* **3**: 79–81.
- Houle D. 1992.** Comparing evolvability and variability of quantitative traits. *Genetics* **130**: 195–204.
- Humphries S, Evans JP, Simmons LW. 2008.** Sperm competition: linking form to function. *BMC Evolutionary Biology* **8**: 319.
- Jamieson IG, Colgan PW. 1992.** Sneak spawning and egg stealing by male threespine sticklebacks. *Canadian Journal of Zoology* **70**: 963–967.
- Kaidbey K, Agin P, Sayre R, Kligman A. 1979.** Photoprotection by melanin – a comparison of black and Caucasian skin. *Journal of the American Academy of Dermatology* **1**: 249–260.
- Keane B, Creel SR, Waser PM. 1996.** No evidence of inbreeding avoidance or inbreeding depression in a social carnivore. *Behavioral Ecology* **7**: 480–489.
- Keller LF, Arcese P, Smith JNM, Hochachka WM, Stearns SC. 1994.** Selection against inbred song sparrows during a natural population bottleneck. *Nature* **372**: 356–357.

- Keller LF, Waller DM. 2002.** Inbreeding effects in wild populations. *Trends in Ecology & Evolution* **17**: 230–241.
- Kemper JH. 1995.** Role of the three-spined stickleback *Gasterosteus aculeatus* L. in the food ecology of the spoonbill *Platalea leucorodia*. *Behaviour* **132**: 1285–1299.
- Kokko H, Ots I. 2006.** When not to avoid inbreeding. *Evolution* **60**: 467–475.
- Kraak SBM, Bakker TCM, Mundwiler B. 1999.** Sexual selection in sticklebacks in the field: correlates of reproductive, mating, and paternal success. *Behavioral Ecology* **10**: 696–706.
- Kvarnemo C, Simmons LW. 2004.** Testes investment and spawning mode in pipefishes and seahorses (Syngnathidae). *Biological Journal of the Linnean Society* **83**: 369–376.
- Largiadèr CR, Fries V, Bakker TCM. 2001.** Genetic analysis of sneaking and egg-thievery in a natural population of the three-spined stickleback (*Gasterosteus aculeatus* L.). *Heredity* **86**: 459–468.
- Leary R, Allendorf F. 1989.** Fluctuating asymmetry as an indicator of stress: implications for conservation biology. *Trends in Ecology and Evolution* **4**: 214–217.
- Louiz I, Ben-Attia M, Ben-Hassine O. 2009.** Gonadosomatic index and gonad histopathology of *Gobius niger* (Gobiidae, Teleost) from Bizerta lagoon (Tunisia): evidence of reproduction disturbance. *Fisheries Research* **100**: 266–273.
- Margulis SW, Walsh A. 2002.** The effects of inbreeding on testicular sperm concentration in *Peromyscus polionotus*. *Reproduction Fertility and Development* **14**: 63–67.
- Mazzi D, Künzler R, Largiadèr CR, Bakker TCM. 2004.** Inbreeding affects female preference for symmetry in computer-animated sticklebacks. *Behavior Genetics* **34**: 417–424.
- Mazzi D, Largiadèr CR, Bakker TCM. 2002.** Inbreeding and developmental stability in three-spined sticklebacks (*Gasterosteus aculeatus* L.). *Heredity* **89**: 293–299.
- Meagher S, Penn DJ, Potts WK. 2000.** Male-male competition magnifies inbreeding depression in wild house mice. *Proceedings of the National Academy of Sciences of the United States of America* **97**: 3324–3329.
- Mehlis M, Bakker TCM, Frommen JG. 2008.** Smells like sib spirit: kin recognition in three-spined sticklebacks (*Gasterosteus aculeatus*) is mediated by olfactory cues. *Animal Cognition* **11**: 643–650.
- Mehlis M, Bakker TCM, Langen K, Frommen JG. 2009.** Cain and Abel reloaded? Kin recognition and male-male aggression in three-spined sticklebacks *Gasterosteus aculeatus* L. *Journal of Fish Biology* **75**: 2154–2162.
- Mortimer D. 1994.** Preparation of human spermatozoa for electron microscopy. In: Mortimer D, ed. *Practical laboratory andrology*. Oxford: Oxford University Press, 168–173.
- Murray R, Kozłowska J, Cutter A. 2011.** Heritable determinants of male fertilization success in the nematode *Caenorhabditis elegans*. *BMC Evolutionary Biology* **11**: 99.
- Ólafsdóttir GÁ, Snorrason SS, Ritchie MG. 2007.** Post-glacial intra-lacustrine divergence of Icelandic threespine stickleback morphs in three neovolcanic lakes. *Journal of Evolutionary Biology* **20**: 1870–1881.
- Parker GA. 1990.** Sperm competition games: sneaks and extra-pair copulations. *Proceedings of the Royal Society of London Series B, Biological Sciences* **242**: 127–133.
- Peer K, Taborsky M. 2005.** Outbreeding depression, but no inbreeding depression in haplodiploid ambrosia beetles with regular sibling mating. *Evolution* **59**: 317–323.
- Plonka P, Passeron T, Brenner M, Tobin D, Shibahara S, Thomas A. 2009.** What are melanocytes really doing all day long . . . ? *Experimental Dermatology* **18**: 799–819.
- Poole A, Lawton C. 2009.** The translocation and post release settlement of red squirrels *Sciurus vulgaris* to a previously uninhabited woodland. *Biodiversity and Conservation* **18**: 3205–3218.
- Postma E, Martini L, Martini P. 2010.** Inbred women in a small and isolated Swiss village have fewer children. *Journal of Evolutionary Biology* **23**: 1468–1474.
- Raeymaekers J, Maes G, Audenaert E, Volckaert F. 2005.** Detecting Holocene divergence in the anadromous-freshwater three-spined stickleback (*Gasterosteus aculeatus*) system. *Molecular Ecology* **14**: 1001–1014.
- Reznick D, Ghalambor C, Nunney L. 2002.** The evolution of senescence in fish. *Mechanisms of Ageing and Development* **123**: 773–789.
- Richards CM. 2000.** Inbreeding depression and genetic rescue in a plant metapopulation. *American Naturalist* **155**: 383–394.
- Rico C, Kuhnlein U, Fitzgerald GJ. 1992.** Male reproductive tactics in the threespine stickleback – an evaluation by DNA fingerprinting. *Molecular Ecology* **1**: 79–87.
- Rioux-Paquette E, Festa-Bianchet M, Coltman D. 2011.** Sex-differential effects of inbreeding on overwinter survival, birth date and mass of bighorn lambs. *Journal of Evolutionary Biology* **24**: 121–131.
- Roldan E, Cassinello J, Abaigar T, Gomendio M. 1998.** Inbreeding, fluctuating asymmetry, and ejaculate quality in an endangered ungulate. *Proceedings of the Royal Society of London Series B, Biological Sciences* **265**: 243–248.
- Saccheri I, Lloyd H, Helyar S, Brakefield P. 2005.** Inbreeding uncovers fundamental differences in the genetic load affecting male and female fertility in a butterfly. *Proceedings of the Royal Society of London Series B, Biological Sciences* **272**: 39–46.
- Savolainen O, Hedrick P. 1996.** Heterozygosity and fitness: no association in Scots pine. *Genetics* **144**: 755–766.
- Sheridan L, Pomiankowski A. 1997.** Fluctuating asymmetry, spot asymmetry and inbreeding depression in the sexual coloration of male guppy fish. *Heredity* **79**: 515–523.
- Sopinka NM, Fitzpatrick JL, Taves JE, Ikonomou MG, Marsh Rollo SE, Balshine S. 2012.** Does proximity to aquatic pollution affect reproductive traits in a wild-caught intertidal fish? *Journal of Fish Biology* **80**: 2374–2383.
- Svensson PA, Forsgren E, Amundsen T, Skold HN. 2005.** Chromatic interaction between egg pigmentation and skin chromatophores in the nuptial coloration of female two-spotted gobies. *Journal of Experimental Biology* **208**: 4391–4397.
- Thünken T, Bakker TCM, Baldauf SA, Kullmann H. 2007.** Active inbreeding in a cichlid fish and its adaptive significance. *Current Biology* **17**: 225–229.

- de Vlaming V, Grossman GD, Chapman F. 1982.** On the use of the gonosomatic index. *Comparative Biochemistry and Physiology* **73**: 31–39.
- Vlieger L, Candolin U. 2009.** How not to be seen: does eutrophication influence three-spined stickleback *Gasterosteus aculeatus* sneaking behaviour? *Journal of Fish Biology* **75**: 2163–2174.
- Waldman B, McKinnon JS. 1993.** Inbreeding and outbreeding in fishes, amphibians and reptiles. In: Thornhill NW, ed. *The natural history of inbreeding and outbreeding: theoretical and empirical perspectives*. Chicago, IL: University of Chicago Press, 250–282.
- Wund MA, Baker JA, Clancy B, Golub JL, Foster SA. 2008.** A test of the ‘flexible stem’ model of evolution: ancestral plasticity, genetic accommodation, and morphological divergence in the threespine stickleback radiation. *American Naturalist* **172**: 449–462.
- Zajitschek SRK, Brooks RC. 2010.** Inbreeding depression in male traits and preference for outbred males in *Poecilia reticulata*. *Behavioral Ecology* **21**: 884–891.
- Zbinden M, Largiadèr CR, Bakker TCM. 2001.** Sperm allocation in the three-spined stickleback. *Journal of Fish Biology* **59**: 1287–1297.
- Zbinden M, Largiadèr CR, Bakker TCM. 2004.** Body size of virtual rivals affects ejaculate size in sticklebacks. *Behavioral Ecology* **15**: 137–140.
- Zbinden M, Mazzi D, Künzler R, Largiadèr CR, Bakker TCM. 2003.** Courting virtual rivals increase ejaculate size in sticklebacks (*Gasterosteus aculeatus*). *Behavioral Ecology and Sociobiology* **54**: 205–209.
- Zieri R, Taboga S, De Oliveira C. 2007.** Melanocytes in the testes of *Eupemphix nattereri* (Anura, Leiuperidae): histological, stereological, and ultrastructural aspects. *Anatomical Record* **290**: 795–800.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Table S1. Descriptive statistics (median, first, and third quartile) of male body measurements (before storage in 70% ethanol), including the values for the duration of nest building and the time that males cared for a nest (days).

Table S2. Table of linear models (‘lm’). In all models, standard length was additionally included as an explanatory variable and never removed to control for possible age influences. 1. ‘lm’ to compare different testis and sperm traits between breeding regime [outbred versus inbred and outbred versus various levels of inbreeding (first and second + third-generation inbred)]; 2. ‘lm’ including interactions between different testis respectively sperm traits and breeding regime.

Appendix S1. Preparing of stickleback sperm for scanning electron microscopy, modified *sensu* Mortimer (1994).

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.